

Infopest verifies that this label is consistent with the NRA
Approved Text Label of 25/11/99

DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SUPRACIDE[®] 400
EMULSIFIABLE CONCENTRATE INSECTICIDE

*Active Constituent: 400 g/L METHIDATHION
(an anticholinesterase compound)*

GROUP 1 B INSECTICIDE

*Controls certain Insect Pests in Orchards, Sub-tropical crops,
Vegetables, Cereals, Pastures, Forage crops, Cotton, Sunflowers,
and other crops as per Directions for Use Table.*

Important: Read the attached booklet before using this product.

1, 5, 10, 20 LITRES

Novartis Crop Protection Australasia Pty. Limited
140-150 Bungaree Road, Pendle Hill NSW 2145

NX

NRA Approval No.: 33041/1199

 **NOVARTIS**

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Very dangerous, particularly the concentrate. Product is poisonous if absorbed by skin contact, inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. DO NOT inhale vapour. Obtain an emergency supply of atropine tablets 0.6 mg. When opening container, preparing spray or using the prepared spray, wear:

- protective waterproof clothing,
- elbow length PVC gloves, and
- full facepiece respirator with combined dust and gas cartridge (canister).

If product or spray on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, respirator (and if rubber wash with detergent and warm water) and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126. If swallowed, give one atropine tablet every 5 minutes until dryness of mouth occurs - if poisoned by skin absorption or through lungs, remove any contaminated clothing, wash skin thoroughly and give atropine tablets as above. Get to a doctor or hospital quickly.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required, refer to the Material Safety Data Sheet.
For a copy phone 1800 025 931.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY.

Novartis has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Novartis accepts no liability for any loss or damage arising from incorrect storage, handling or use.

® Registered Trademark of Novartis AG, Basel, Switzerland.

<p>UN No. 3018 ORGANOPHOSPHORUS PESTICIDE, LIQUID, TOXIC, N.O.S. (CONTAINS METHIDATHION) Packaging Group II HAZCHEM 2X</p>		<p>In a Transport Emergency Dial 000, Police or Fire Brigade.</p> <p>For specialist advice in an emergency only call 1 800 033 111 (24 hours)</p>
---	---	---

BAR CODE

Batch Number	
Date of Manufacture	

DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SUPRACIDE[®] 400
EMULSIFIABLE CONCENTRATE INSECTICIDE

*Active Constituent: 400 g/L METHIDATHION
(an anticholinesterase compound)*

*Controls certain Insect Pests in Orchards, Sub-tropical crops,
Vegetables, Cereals, Pastures, Forage crops, Cotton, Sunflowers,
and other crops as per Directions for Use Table.*

Important: Read the attached booklet before using this product.

1, 5, 10, 20 LITRES

Novartis Crop Protection Australasia Pty. Limited
140-150 Bungaree Road, Pendle Hill NSW 2145

NX

NRA Approval No.: 33041/1199

 **NOVARTIS**

Table of Contents**Page**

Apples	
Apricots	
Avocados	
Beans	
Broccoli	
Brussels Sprouts	
Cabbage	
Cauliflower	
Cereals	
Citrus	
Coffee	
Cotton	
Custard Apples	
Eggplants	
Flower Seedlings	
Forage Crops	
Garlic	
Grapes	
Kale	
Kohlrabi	
Lettuce	
Lucerne	
Lupins	
Macadamia Nuts	
Mangoes	
Oilseeds	
Onion	
Orchids	
Ornamentals	
Passionfruit	
Pasture	
Pears	
Peas	
Potatoes	
Rice	
Stone Fruit	
Sunflower	
Tomatoes	
Trees and shrubs	
Turf	
Vegetable Seedlings	

DIRECTIONS FOR USE

Restraint: The effect of this product could be diminished if rain falls within 6 hours of spraying.

ORCHARDS AND VINEYARDS

Crop	Insect	State	Rate		WHP	Critical Comments
			High Volume (rate/100 L)	Aircraft (rate/ha)		
Apples, Pears Spring and Summer Treatment	Codling Moth, Lightbrown Apple Moth, Two-spotted Mite	Qld only	125 mL	-	14 days	Start application at petal fall and continue spraying at 21 day intervals as required. For best results ensure thorough coverage of foliage and fruit. Refer to "Precaution" in General Instructions.
	Codling Moth, Lightbrown Apple Moth, Two-spotted Mite, Woolly Aphid	NSW, Vic, Tas, SA, WA only				Start application at petal fall and continue spraying at 21 day intervals as required. For best control of San José scale and mealybug, at least 3 applications commencing at petal fall are recommended. If codling moth and lightbrown apple moth are severe problems, shorten interval to 14 days between treatments until pest activity decreases. Spraying interval should be shortened if heavy rain falls soon after application. For best results ensure thorough coverage of foliage and fruit. Where woolly aphid, mealybug and San José scale are present, pay particular attention to thorough coverage of limbs and branches. These applications will keep woolly aphid in check during the period they are applied. The addition of a specific miticide will be necessary when mite resistance occurs. Frequent observations of mites are desirable to avoid population build up.
	San José Scale	Qld, NSW, Vic, SA, WA only				For control of San José scale apply as late as possible during the dormant period, preferably at budswell. Ensure thorough coverage of limbs and branches. If desired, 1-2 L of dormant oil may be added to the spray mixture, but is not necessary for good control of San José scale. DO NOT apply to stone fruit after the dormant period as partial defoliation of some varieties may occur.
	Mealybug	Vic, Tas, SA, WA only				
Apples, Pears, Stone Fruit Dormant Treatment	San José Scale	Qld, NSW, Vic, SA only WA only	125 mL with 3 L dormant spraying oil.			
Apricots Dormant Treatment	Grapevine Scale	Qld, WA only	125 mL			
Avocados	Banana-spotting Bug, Fruitspotting Bug, Monolepta Beetle				7 days	Spray when pests are active. Repeat applications at 7-10 day intervals.
	Latania Scale, Redbanded Thrips					Spray thoroughly as required.

Crop	Insect	State	Rate		WHP	Critical Comments	
			High Volume (rate/100 L)	Aircraft (rate/ha)			
Grapes Dormant Treatment	Grapevine Scale	Qld, NSW, Vic, SA, WA only	125 mL	-	14 days	Apply during winter months prior to bud movement.	
Grapes Spring/Summer Treatment	Grapevine Moth	Qld, NSW, SA, WA only	65 mL	650 mL	14 days	Spray when caterpillars are noticed during spring and early summer. Repeat application if necessary.	
	Grapevine Scale	Vic only	75 mL	750 mL		Spray during late Nov-Dec. For late table grapes, spray early Nov and again late Dec. Thorough coverage is essential.	
	Longtailed Mealybug	Qld, NSW, SA, WA only	125 mL	1.3 L		-	Spray when caterpillars are first noticed and repeat applications if necessary at 14-21 day intervals.
		Vic only	-				
	Lightbrown Apple Moth	Qld, NSW, SA, WA only	-	1.3 L		-	Spray during spring and preferably at monthly intervals or at first signs of adult feeding or egg laying activity.
		Vic only	-				
Common Auger Beetle, Elephant Weevil, Fig Longicorn, Vine Weevil	NSW, WA only	-	-	-			
Citrus	Mealybug	SA, WA only	125 mL with a non ionic surfactant at label rates (Qld, Vic, SA, WA); 125 mL (NSW); OR 125 mL with 1 L miscible summer spraying oil. DO NOT add oil under very hot conditions	-	21 days	Apply during Nov-Feb period. Two sprays may be required under conditions of very heavy red scale infestation, the first during late Nov/Dec, the second during Jan/Feb. In SA spraying may commence one month earlier and oil should not be added when spraying in Nov/ Dec. Ensure thorough coverage of foliage, limbs and branches.	
	Red Scale	NSW, Vic, SA, WA only				Spray when larvae are on the leaves - mid-Dec in warmer districts, to mid-Jan in cooler areas of NSW. Spray thoroughly.	
	White Wax Scale	NSW only				Spray in Dec/Jan period. Apply a second spray 2 weeks later if infestation very heavy. Spray thoroughly.	
		WA only				Spray in late Nov/early Dec and again in late Jan/early Feb. Ensure thorough coverage of foliage, limbs and branches. DO NOT use oil spray mixture on Glen Retreat Mandarins. Scale spray applications also control orange fruit borer, mealybug and black citrus aphid. DO NOT apply more than twice in any one season. Refer to "Precaution" in General instructions.	
	Red Scale, White Wax Scale	Qld only				Apply as necessary at growth flushes.	
	Circular Black Scale, Pink Wax Scale, Soft Brown Scale	Qld, WA only					
	Citrus Leafminer					125 mL with a non ionic surfactant at label rates	

Crop	Insect	State	Rate		WHP	Critical Comments
			High Volume (rate/100 L)	Aircraft (rate/ha)		
Citrus	Bronze Orange Bug, Spined Citrus Bug	Qld, WA only	125 mL with a non ionic surfactant at label rates	-	21 days	Use as required. Apply before bronze orange bugs reach adult stage.
			30 mL with a non ionic surfactant at label rates			Use where integrated control practices are being followed. These low rates of SUPRACIDE may not provide equivalent control to full strength recommendations, but will suppress aphid and bug populations, whilst having minimum effect on desirable insects such as parasites of scale insects and predatory ladybirds.
	Black Citrus Aphid		25 mL			Apply at any time of year, directing the spray to achieve thorough coverage of trunk and limbs. For severe infestation the canopy of the tree should also be thoroughly sprayed. The addition of oil as for other insects may aid in control but is not necessary for white louse scale control.
	White Louse Scale		125 mL with a non ionic surfactant at label rates			Monitor fruit early Feb to harvest on a fortnightly basis. Spray if 10 or more freshly fallen fruit per tree are infested at any one time.
	<i>Blastobasis</i> spp., Sorghum Head Caterpillar		125 mL			Check fruit in Oct/Nov. Spray if more than 1 in 10 fruit stalks are infested.
	Hemispherical Scale, Long Soft Scale, Purple Scale					Check leaves in Oct/Nov. Spray if more than 1 in 50 leaves are infested.
	Green Coffee Scale, Pulvinaria Scale					Spray if more than 1 in 3 twigs galled. Apply early to mid-Dec. Timing is critical.
	Citrus Gall Wasp					

SUB-TROPICAL CROPS

Crop	Insect	State	Rate	WHP	Critical Comments
			High Volume (rate/100 L)		
Coffee (non-bearing)	Mealybug	Qld, WA only	125 mL with 1 L white oil	21 days	Apply to established infestations.
Custard Apples	Fruitspotting Bug	Qld, NSW, WA only	65 mL	7 days	Apply when pest active.
	Black Scale, Long Soft Scale, Soft Brown Scale		125 mL with a non ionic surfactant at label rates		Apply thoroughly as required.
	Mealybug				Apply approximately 1 month before harvest and repeat 2-3 weeks later. Good spray coverage of fruit is essential.

Crop	Insect	State	Rate	WHP	Critical Comments	
			High Volume (rate/100 L)			
Macadamia Nuts	Macadamia Felted Coccid	Qld, NSW, WA only	125 mL	21 days	Spray when pests noticed. Repeat applications monthly as necessary. Spray thoroughly.	
	Macadamia Nutborer, Macadamia Twig-girdler				Spray as required.	
	Black Citrus Aphid, Banana-spotting Bug, Macadamia Leafminer	Qld, WA only				
	Fruitspotting Bug	Qld, NSW, WA only	65 mL			Spray when pests active.
	Black Scale, Pink Wax Scale, Soft Brown Scale, White Wax Scale	Qld, WA only	125 mL with a non ionic surfactant at label rates			Spray thoroughly in late Nov/early Dec, when scale damage is noticed on branches of trees.
	Long Soft Scale, Macadamia Mussel Scale, Macadamia White Scale	Qld, NSW, WA only				
	Citrus Mealybug, Hibiscus Mealybug, Longtailed Mealybug	Qld, WA only				Spray when pests or honey dew are noticed.
Mangoes	Mango Scale				Spray thoroughly in late Nov/ early Dec.	
	Pink Wax Scale	Qld, NSW, WA only	125 mL with 1L of a water miscible oil		Spray as required.	
	Mango Tipborer					
	Leafminers - (<i>Acrocercops</i> spp.)	Qld, WA only			Spray when pests are active.	
	Mango Seed Weevil	Qld, NSW, WA only			Apply as a cover spray Feb-Mar when trees are flushing and beetles emerge from seeds. A repeat canopy spray when fruit is 2-3 cm in diameter will kill the adult weevil.	
Passionfruit	Mealybug		125 mL with a non ionic surfactant at label rates	14 days	Apply 1-2 applications (2-3 weeks apart) depending on the degree of infestation.	
	Red Scale				Apply early Dec and again late Jan/early Feb.	
	Scale Insects		125 mL		Apply the spray when scale populations build up to high levels. Ensure thorough coverage. Better activity against black scale may be achieved by application when the insects are in the young stages.	

SUNDRY CROPS AND PASTURES

Crop	Insect	State	Rate		WHP	Critical Comments
			By Boom Spray, Aircraft, Mister (rate/ ha)	High Volume (rate/100 L)		
Lupins	Cowpea Aphid	SA, WA only	1 L	-	7 days	Apply when pests active.
	Lucerne Flea, Redlegged Earth Mite	Vic, SA, WA only	90 mL		1 day	Spray when pests appear in large numbers at seedling emergence.
Cereals	Blue Oat Mite	All States				Spray when pests appear in large numbers, 3-6 weeks after autumn rains.
Cereals, Pasture and Forage Crops	Lucerne Flea, Redlegged Earth Mite	NSW, Vic, Tas, SA, WA only				If pest numbers build up during the season, spray again. For best results, avoid spraying when pests are sheltering. Spray when at least 2.5 cm cover of pasture or crop is present.
	Common Armyworm	Qld, NSW, Vic, SA, WA only	1.4 L		7 days for grazing/ fodder OR 6 weeks for grain for human consumption	Spray at first sign of infestation.
	Southern Armyworm (Barley Grub)	NSW, Vic, Tas, SA, WA only				
Cereals, Pastures, Forage Crops, Oilseeds, Lucerne Bare earth situation	Blue Oat Mite, Lucerne Flea, Redlegged Earth Mite	NSW, Vic, SA, WA only	200 mL		7 days	Apply as a ground spray, immediately prior to seedling emergence using sufficient water to give good coverage. If pest activity is severe, also spray the headlands and surrounding vegetation prior to emergence.
Pasture	Blackheaded Pasture Cockchafer	NSW, Vic, Tas, SA, WA only	1 L		7 days for grazing/ fodder OR 6 weeks for grain for human consumption	Spray when rain is imminent. Rain is required for cockchafer to surface.
Lucerne	Lucerne Leafroller	All States	850 mL			Spray when the pest first appears and repeat as necessary.
	Sitona Weevil	NSW, Vic, SA, WA only	1.4 L			Apply at first sign of serious infestation. Re-spraying may be necessary if conditions favour further emergence of weevils from the soil.
	Spotted Alfalfa Aphid	Qld, NSW, SA, WA only	750 mL			Apply as soon as practical following cutting and repeat as necessary.
	Bluegreen Aphid	Qld, NSW, WA only				
Rice	Common Armyworm	Qld, WA only	1.4 L		6 weeks	Spray at first sign of infestation.

Crop	Insect	State	Rate		WHP	Critical Comments
			By Boom Spray, Aircraft, Mister (rate/ ha)	High Volume (rate/100 L)		
Cotton	Mites (Knockdown and short residual control)	Qld, NSW only	1.4 L	-	3 days	Ideally spray when infestation is light to moderate. If mite populations are high or conditions favour a build-up, repeat application in 5-7 days. Spray with good coverage in at least 20 L water/ha. DO NOT apply more than 3 times during the season.
		Sunflower	Grey Cluster Bug, Rutherglen Bug	Qld, NSW, Vic, WA only		
Turf	Argentine Stem Weevil, Couchtip Maggot	SA, WA only	-	500 mL (8 L spray/ 100 square metres)	Nil	Apply at 10 day intervals as required. DO NOT graze treated turf or feed grass clippings from treated turf to animals.
	Couchgrass Mite, Couchgrass Scale	Qld, SA, WA only		750 mL (14 L spray/ 100 square metres)		
	Mealybug					

VEGETABLES

Crop	Insect	State	Rate			WHP	Critical Comments
			Boom Spray (rate/ha)	High Volume (rate/ 100L)	Knapsack (rate/15L)		
Beans	Aphids	Qld, Vic, Tas, SA, WA only	1.4 L	125 mL	20 mL	7 days	Apply at the first signs of infestation. Re-spray as necessary.
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale, Kohlrabi	Cabbage Aphid, Cabbage Moth, Cabbage White Butterfly, Green Peach Aphid	All States					Apply at the first sign of activity. Continue applications every 10-14 days while pests are active, with intervals of 14 days in Qld. Thorough coverage is required on larger plants.
Eggplants	Aphids, Leafhoppers	Qld, Vic, Tas, SA, WA only					Spray when necessary.
Lettuce	Aphids	SA, WA only					Apply at the first signs of infestation. Re-spray as necessary.
Onions	Onion Thrips	All States	750 mL	75 mL	10 mL		Spray thoroughly when pests observed. A follow-up application 7- 10 days later may be necessary.
Garlic		Qld, Vic, Tas, SA, WA only					
Peas	Aphids, Loopers	WA only	1.4 L	125 mL	20 mL		Apply at the first signs of infestation. Re-spray as necessary.

Crop	Insect	State	Rate			WHP	Critical Comments
			Boom Spray (rate/ha)	High Volume (rate/100 L)	Knapsack (rate/15 L)		
Potatoes	Rutherglen Bug	All States	1 L	-	-	7 days	Spray at the first of infestation.
	Loopers	Qld, Vic, Tas, SA, WA only	1.4 L	125 mL	20 mL		Repeat applications as necessary.
Tomatoes	Leafhoppers	All States					Spray when necessary.
	Aphids	All States					Spray at first sign of infestation. Repeat applications at 10-14 day intervals or as required.
	Cutworms	Qld, Vic, Tas, SA, WA only	1.4 to 2.8 L				Spray thoroughly, including soil around plants.
	Onion Thrips	Qld, Vic, Tas, SA, WA only	750 mL	75 mL	10 mL		Spray thoroughly when pests observed. Re-spray as necessary.
	Rutherglen Bug	All States	1 L	-	-		Spray during periods of serious infestation. Repeat application as necessary if reinfestation occurs from surrounding areas.

ORNAMENTALS, TREES, SHRUBS, ANNUALS

Plant	Insect	State	Rate	WHP	Critical Comments
			High Volume (rate/100 L)		
Orchids	Aphids, Black Scale, Orchid Beetle, Orchid Scale, Soft Brown Scale, Slugs and Snails, Thrips	Qld, NSW, Vic, Tas, SA, WA only	125 mL	Nil	Apply spray thoroughly at the first signs of pest activity. Scale insects are most effectively controlled when sprayed in the crawler (larval) stage. DO NOT spray during flowering period.
Trees and Shrubs in nurseries, parks and forestry situations	Aphids, Armyworm, Cutworm, Leafhoppers, Leafminers, Lerp Insects, Sawflies, Scale Insects, Soft Scales, Thrips				Apply at the first sign of infestation and repeat if re-infestation occurs. Spray scale insects in the crawler stage for most effective control.
Flower and Vegetable Seedlings	Aphids, Armyworm, Cutworm, Leafhoppers, Thrips, Weevils		100 to 125 mL		Apply at the first sign of infestation and repeat at 10-14 day intervals as necessary. Some leaf scorching may occur on tender young foliage.
Ornamentals in nurseries, parklands and gardens.	Aphids, Armyworm, Cutworm, Grasshoppers, Leafhoppers, Mealybug, Scale Insects, Thrips, Weevils				Apply at the first sign of infestation and repeat at 10-14 day intervals.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN MARKET.

IN TASMANIA THIS PRODUCT MUST NOT BE APPLIED BY AIRCRAFT WITHOUT THE SPECIFIC APPROVAL OF THE REGISTRAR OF PESTICIDES.

WITHHOLDING PERIODS:**Vegetables, Custard Apples:**

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

Apples, Pears, Stone Fruits, Grapes, Passionfruit:

DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

Citrus, Macadamia Nuts, Mangoes:

DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

Oil Seed Crops (including Cotton):

DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

Pasture and Forage Crops (including Lucerne, Lupins, Sorghum and Cereal Crops):

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION OF 90mL PRODUCT PER HECTARE.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION IN EXCESS OF 90 mL PER HECTARE.

Grain Crops (including Grain Sorghum, Rice and Cereal Crops):

DO NOT APPLY TO SORGHUM AND CEREAL CROPS WITHIN 6 WEEKS OF HARVEST, IF GRAIN INTENDED FOR HUMAN CONSUMPTION.

GENERAL INSTRUCTIONS**Mixing**

Fill the spray tank $\frac{1}{4}$ full of water. With the agitator or by-pass in operation, slowly add the required amount of product to the spray tank while continuing to fill the tank with water. Add the remainder of the water while continuing to agitate. If oils or surfactants are to be added to the spray tank, add these just before the tank is full of water. Continue agitation through-out the spray operation.

Application

General rates of application are :

- * High volume orchard and power sprayers: 65 to 125 mL/100 L water.
- * Boom sprayer, aircraft, misting machine: 90 mL/ha for earth mites and lucerne flea; 850 mL to 1.4 L/ha for other pests. For aerial application use a min. of 20-30 L water/ha.
- * Knapsack sprayer: 10 to 20 mL per 15 L knapsack.

Addition of surfactant

To improve spray coverage especially on cabbages, cauliflowers and broccoli add a non ionic surfactant at registered label rates.

Insecticide resistance warning GROUP 1 B INSECTICIDE

For insecticide resistance management SUPRACIDE is a Group 1B insecticide. Some naturally occurring insect biotypes resistant to SUPRACIDE and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if SUPRACIDE and other Group 1B insecticides are used repeatedly. The effectiveness of SUPRACIDE on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Novartis Crop Protection accepts no liability for any losses that may result from failure of SUPRACIDE to control resistant insects. SUPRACIDE may be subject to specific resistance management strategies. For further information contact your supplier, Novartis Crop Protection representative or local agricultural department agronomist.

PRECAUTIONS

This product is deleterious to beneficial insects. Care should be exercised when applying into pest management situations, e.g. citrus and deciduous fruits. In Qld, do not apply SUPRACIDE sprays for citrus insect pests more than twice in any one season.

Re-entry Period: DO NOT enter treated areas without protective clothing until 48 hours after spraying.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, ENVIRONMENT AND BEES

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or the used container. Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and

return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Very dangerous, particularly the concentrate. Product is poisonous if absorbed by skin contact, inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. DO NOT inhale vapour. Obtain an emergency supply of atropine tablets 0.6 mg. When opening container, preparing spray or using the prepared spray, wear:

- protective waterproof clothing,
- elbow length PVC gloves, and
- full facepiece respirator with combined dust and gas cartridge (canister).

If product or spray on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, respirator (and if rubber wash with detergent and warm water) and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126. If swallowed, give one atropine tablet every 5 minutes until dryness of mouth occurs - if poisoned by skin absorption or through lungs, remove any contaminated clothing, wash skin thoroughly and give atropine tablets as above. Get to a doctor or hospital quickly.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required, refer to the Material Safety Data Sheet.

For a copy phone 1800 025 931.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY.

Novartis has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Novartis accepts no liability for any loss or damage arising from incorrect storage, handling or use.

® Registered Trademark of Novartis AG, Basel, Switzerland.

* Registered Trademark.